

Title	The relationship between IGF-I and -II concentrations and body composition at birth and over the first 2 months
Authors	Hawkes, Colin P.;Grimberg, Adda;Kenny, Louise C.;Kiely, Mairead E.;Hourihane, Jonathan O'B.;Irvine, Alan D.;McPhaul, Michael J.;Caulfield, Michael P.;Zemel, Babette S.;Murray, Deirdre M.
Publication date	2019-02-11
Original Citation	Hawkes, C. P., Grimberg, A., Kenny, L. C., Kiely, M., Hourihane, J. O. B., Irvine, A. D., McPhaul, M. J., Caulfield, M. P., Zemel, B. S. and Murray, D. M. (2019) 'The relationship between IGF-I and -II concentrations and body composition at birth and over the first 2 months', Pediatric Research, 85(5), pp. 687-692. doi: 10.1038/s41390-019-0331-x
Type of publication	Article (peer-reviewed)
Link to publisher's version	<a href="https://www.nature.com/articles/s41390-019-0331-x">https://www.nature.com/articles/s41390-019-0331-x</a> - 10.1038/s41390-019-0331-x
Rights	© 2018 the authors. Published by Springer Nature. This is a post-peer-review, pre-copyedit version of an article published in Pediatric Research. The final authenticated version is available online at: <a href="http://dx.doi.org/10.1038/s41390-019-0305-z">http://dx.doi.org/10.1038/s41390-019-0305-z</a>
Download date	2023-05-05 04:52:20
Item downloaded from	<a href="http://hdl.handle.net/10468/7772">http://hdl.handle.net/10468/7772</a>



# UCC

**University College Cork, Ireland**  
Coláiste na hOllscoile Corcaigh

## The relationship between IGF-I and -II concentrations and body composition at birth and over the first two months

**Cite this article as:** Colin P Hawkes, Adda Grimberg, Louise C Kenny, Mairead Kiely, Jonathan O. B Hourihane, Alan D Irvine, Michael J. McPhaul, Michael P. Caulfield, Babette S Zemel and Deirdre M Murray, The relationship between IGF-I and -II concentrations and body composition at birth and over the first two months, *Pediatric Research* doi:10.1038/s41390-019-0331-x

This Author Accepted Manuscript is a PDF file of an unedited peer-reviewed manuscript that has been accepted for publication but has not been copyedited or corrected. The official version of record that is published in the journal is kept up to date and so may therefore differ from this version.

Terms of use and reuse: academic research for non-commercial purposes, see here for full terms. <https://www.nature.com/authors/policies/license.html#AAMtermsV1>

Author accepted manuscript

## The relationship between IGF-I and –II concentrations and body composition at birth and over the first two months

\*Colin P Hawkes<sup>1,2,3</sup>, Adda Grimberg<sup>1,3</sup>, Louise C Kenny<sup>5,6</sup>, Mairead Kiely<sup>5,7</sup>, Jonathan O'B Hourihane<sup>4,5</sup>, Alan D Irvine<sup>8</sup>, Michael J. McPhaul<sup>9</sup>, Michael P. Caulfield<sup>9</sup>, Babette S Zemel<sup>3,10</sup>, Deirdre M Murray<sup>4,5</sup>.

<sup>1</sup>Division of Endocrinology and Diabetes, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania.

<sup>2</sup>The National Children's Research Centre, Dublin, Ireland.

<sup>3</sup>Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, Pennsylvania.

<sup>4</sup>Department of Paediatrics and Child Health, University College Cork, Cork, Ireland.

<sup>5</sup>The Irish Centre for Fetal and Neonatal Translational Research, Ireland.

<sup>6</sup>Faculty of Health and Life Sciences, University of Liverpool, United Kingdom.

<sup>7</sup>School of Food and Nutritional Science, University College Cork, Ireland.

<sup>8</sup>Department of Clinical Medicine, Trinity College Dublin, Ireland.

<sup>9</sup>Quest Diagnostics Nichols Institute, San Juan Capistrano, California

<sup>10</sup>Division of Gastroenterology, Hepatology and Nutrition, The Children's Hospital of Philadelphia, Philadelphia, Pennsylvania.

### Correspondence should be addressed to

Colin Hawkes, MD PhD, Division of Endocrinology and Diabetes, The Children's Hospital of Philadelphia, Suite 11NW30, 3401 Civic Center Blvd, Philadelphia, PA, 19104.

Email: [hawkesc@email.chop.edu](mailto:hawkesc@email.chop.edu)

Tel: 01-215-590-3174

Fax: 01-215-590-3053

### Category of Study: Clinical Research

**Statement of Financial Support:** Dr. Hawkes is supported by a PhD grant by the National Children's Research Centre, Dublin, Ireland and Dr. Grimberg was supported by grant 1R01 HD57037 from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD). The SCOPE Ireland Study was funded by the Health Research Board of Ireland (CSA 02/2007). L.C.K. is a Science Foundation Ireland (SFI) principal investigator (PI) (08/IN.1/B2083). L.C.K., D.M. and M.K. are PIs in the SFI-funded Research Centre, INFANT (12/RC/2272). The Cork BASELINE birth cohort is funded by the National Children's Research Centre.

**Disclosures Statement:** AG serves on the Steering Committee of the Pfizer International Growth Study Database. MJM and MPC are employees of Quest Diagnostics. All other authors have no financial relationships relevant to this article to disclose.

**Acknowledgement:** The authors would like to acknowledge Julia Larsen at Quest Diagnostics and Klaudia Sikora at University College Cork for sample handling and data retrieval.

## Abstract

**Background:** Insulin-like growth factors (IGF)-I and -II play an important role in prenatal growth. During the first two months from birth, body fat doubles, and rapid weight gain during this time increases future risk of cardiometabolic disease.

**Objective:** To determine if IGF measurements at birth associate with body composition and the trajectory of its changes in the first two months.

**Methods:** Umbilical cord IGF-I and -II concentrations were measured in term infants. Air displacement plethysmography was performed at birth and two months. Fat mass (FM) and fat free mass (FFM) were corrected for infant length (L) to  $FM/L^3$  and  $FFM/L^2$  respectively.

**Results:** In 601 (317 male) infants, IGF-I concentrations at birth were associated with  $FM/L^3$  and  $FFM/L^2$  Z-scores at birth ( $R^2=0.05$  and  $0.04$  respectively,  $P<0.001$ ), and IGF-II concentrations were associated with  $FFM/L^2$  Z-scores at birth ( $R^2=0.01$ ,  $P=0.02$ ). Lower IGF-I concentrations were weakly associated with increases in  $FM/L^3$  Z-scores over the first two months ( $R^2=0.01$ ,  $P=0.003$ ).

**Conclusion:** IGF-I concentrations at birth are associated with adiposity and lean mass at birth and inversely with the trajectory of FM accumulation over the first two months. IGF-I measurements only account for a small amount of the variance in these measures.

## Introduction

The *in utero* environment and early infant growth play an important role in the development of adult disease. Low birth weight is associated with increased risk of cardiovascular disease in adulthood(1) and offspring of mothers exposed to famine during pregnancy have shown an increased risk of obstructive airway disease(2), metabolic syndrome(3), and psychiatric disease(4, 5). Following birth, rapid early weight gain also adversely affects risk of obesity(6), insulin resistance(7), premature adrenarche(8) and cardiovascular disease(1, 6). These early-life weight changes may have additional implications for future health, as the obese phenotype often tracks from early childhood through adulthood(9).

Insulin-like growth factor (IGF)-I and -II play an important role in regulating fetal growth. Infants with IGF-I and -II signaling defects have significant prenatal growth failure(10). Regulation of IGF-I and -II *in utero* appears to be more dependent on nutrition(11) than growth hormone (GH). Infants with GH deficiency have normal size at birth(12) whereas infants with intrauterine growth restriction have reduced concentrations of IGF-I and -II(13, 14). This correlation with weight at birth is stronger for IGF-I than for IGF-II concentrations(14). Animal models also show that expression of IGF-I and -II is reduced in response to intrauterine nutritional deprivation(15).

Birth weight is often used as a proxy for adiposity, but these do not always correlate(16, 17). It is possible that IGF-I and -II concentrations are associated more closely with lean and fat mass than with birth weight. Infants born small for gestational age (SGA) have a greater deficit in

body fat than in lean mass and have lower IGF-I concentrations(18), suggesting that IGF-I levels may be more closely associated with body fat than lean mass at birth. IGF-I measurement may also be useful in assessing early growth trajectories. In children born appropriate for gestational age, higher IGF-I concentrations at three months of age are associated with increased weight gain over the preceding three months(19).

There are limited data describing the association of IGF-I and -II concentrations with detailed body composition measurements at birth, or assessing the predictive value of IGF-I and -II concentrations at birth on subsequent dynamic changes in body composition. The aim of this study is to determine if IGF-I and IGF-II measurements at birth are associated with body composition at birth and with the trajectory of body composition changes in the first two months from birth.

## Methods

### *Subjects*

The Cork BASELINE birth cohort study (ClinicalTrials.gov NCT: 01498965) is a prospective study that recruited infants born in Cork, Ireland between August 2008 and August 2011(20). Children born between 37 and 42 weeks' gestation to low risk first-time mothers were eligible for inclusion in this study. Exclusion criteria were non-singleton pregnancies, known major fetal anomalies, maternal prepregnancy hypertension, diabetes, renal disease, systemic lupus erythematosus, antiphospholipid syndrome, major uterine anomalies, cervical cone biopsy and  $\geq 3$  miscarriages. Ethical approval was granted by the Clinical Research Ethics Committee of the Cork Teaching Hospitals.

The first day of the last menstrual period (LMP) was used to determine gestational age. Ultrasound gestational age was used if performed before 16 weeks' gestation and if there was a discrepancy of more than six days between this and LMP. Ultrasound gestational age was also used if there was a discrepancy of more than ten days between LMP and the ultrasound measurement at 20 weeks' gestation.

### *Body Composition assessment*

Air displacement plethysmography (ADP) (PEAPOD<sup>TM</sup> Infant Body Composition System, Life Measurement Inc, Concord, CA)(21) was used to measure body composition at birth and two months of age. Standardized protocols were used to assess body composition as per manufacturer specifications. The naked infant, without umbilical cord clip, was weighed on a high precision

scale and placed in a closed chamber. Pressure and volume changes in the chamber were used to calculate air displacement, and body density was determined. Age and sex specific FFM density values, and the density of fat were then used to calculate FM and FFM(22, 23).

Birth measurements were included if they occurred within the first four days from birth. In this cohort, 90% of two month measurements were taken between 49 and 86 days of age, and two month measurements were included if they occurred within this window. FM and FFM are dependent on infant size, so these measurements were adjusted for infant length in order to remove the confounding effect of size(24, 25). We have previously shown that  $FM/L^3$  ( $FM/L^3$ ) and  $FFM/L^2$  are the optimal indices for this correction in the first three months from birth(16). Age- and sex-specific Z-scores for  $FM/L^3$  and  $FFM/L^2$  parameters have been determined from this birth cohort, and have been previously described elsewhere(16).

#### *Sample collection and storage*

Umbilical cord samples were collected at birth. They were processed to serum within three hours of collection, and stored at  $-80^{\circ}\text{C}$  until analyzed.

#### *Mass Spectrometry*

IGF-I and -II concentrations were measured using liquid chromatography-mass spectrometry (LCMS) (Quest Diagnostics, Madison, NJ). This methodology is described in detail elsewhere(26, 27). Isotopically labeled IGF-I was added to the sample as an internal standard to correct for any analyte lost in binding protein extraction. IGF-I and -II were separated from their binding proteins by an acid ethanol extraction followed by automated online extraction and



analytical chromatography using an Aria TX-4 (Thermo-Fisher, San Jose, CA). IGF-I and -II were quantified using a time-of-flight mass spectrometer using narrow mass extraction of full scan spectra. Performance characteristics of this assay have been previously described(26, 27). For IGF-I measurement, the interassay coefficient of variation and percent recovery were 5 and 104% at 100 ng/mL, 5.2 and 103% at 400 ng/mL, and 3.5 and 103% at 700 ng/mL. For IGF-II measurement, the interassay coefficient of variation and percent recovery were 6.1 and 102% at 200 ng/mL, 3.2 and 99% at 500 ng/mL, and 5.3 and 99% at 1,200 ng/mL.

### *Statistical Analysis*

Analysis of  $FM/L^3$  and  $FFM/L^2$  also utilized age- and sex-specific Z-scores at birth and two months. The change in Z-score between birth and two months was calculated by subtracting the age- and sex-specific Z-score at birth from the two months Z-score. Gestational age- and sex-specific Z-scores for IGF-I and -II concentrations were also used in this analysis, and the development of these reference data are described elsewhere(14). The LCMS assay cannot detect IGF-I concentrations below 16 ng/ml or IGF-II concentrations below 32 ng/ml. When samples had concentrations below these values, a concentration of 15 ng/ml for IGF-I or 31 ng/ml for IGF-II was assigned.

Linear regression analysis was used to test associations between IGF-1 and -II concentration Z-scores with body composition. When evaluating the relationship between IGF Z-scores at birth and changes in  $FM/L^3$  and  $FFM/L^2$  Z-scores over the first two months, partial correlations were used to correct for  $FM/L^3$  and  $FFM/L^2$  Z-scores at birth.

Data analyses were performed using SPSS 21.0 (IBM, New York). Mean (SD) were reported for normally distributed data and compared using Student's independent sample T-tests. Linear regression analysis was used to evaluate the relationship between continuous variables.

Author accepted manuscript

## Results

There were 2137 infants enrolled in the Cork BASELINE birth cohort. Of these, 105 were excluded from this study due to gestational age less than 37 weeks, 932 did not have sufficient cord blood available for IGF measurement, and a further 499 did not have body composition measurement performed both at birth and two months. Complete measurements of IGF-I and -II at birth and ADP measurements at birth and two months of age were available in 601 term infants (317 male) (Figure 1). Characteristics of these subjects are shown in Table 1. At birth, IGF-I concentration Z-Score correlated positively with IGF-II concentration Z-Score ( $R^2=0.08$ ,  $P < 0.001$ ).

### *1. Association of IGF-I and -II concentrations at birth with body composition at birth*

Increased IGF-I concentrations were associated with higher FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at birth ( $R^2=0.05$ ,  $P < 0.001$  and  $R^2=0.04$ ,  $P=0.016$  respectively) although these accounted for a small amount of the variance in body composition at birth (Figure 2). IGF-II concentrations at birth were associated with FFM/L<sup>2</sup> Z-score at birth ( $R^2=0.01$ ,  $P=0.02$ ) but not with FM/L<sup>3</sup> Z-score at birth ( $R^2=0.002$ ,  $P=0.4$ ).

### *2. Association of IGF-I and -II concentrations at birth with body composition trajectory over the first 2 months*

FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at 2 months were not associated with IGF-I ( $R^2=0.002$ ,  $P=0.3$  and  $R^2=0.001$ ,  $P=0.5$  respectively) or IGF-II ( $R^2=0.004$ ,  $P=0.1$  and  $R^2=0.002$ ,  $P=0.3$  respectively) Z-

scores at birth. This indicates that body composition at two months was not predicted by IGF-I or -II concentrations at birth.

Although IGF-I and -II concentrations at birth were not associated with body composition at two months, there was a weak but significant association with change in body composition over the first two months. Higher IGF-I Z-scores at birth were associated with reductions in FM/L<sup>3</sup> Z-scores ( $R^2=0.05$ ,  $P < 0.001$ ) and FFM/L<sup>2</sup> Z-scores ( $R^2=0.03$ ,  $P < 0.001$ ) between birth and two months of age. However, IGF-II concentrations were not associated with changes in FM/L<sup>3</sup> Z-scores ( $R^2=0.006$ ,  $P=0.05$ ) or FFM/L<sup>2</sup> Z-scores ( $R^2=0.003$ ,  $P=0.02$ ) over the same timeframe (Figure 3).

To control for the potential effect of FM/L<sup>3</sup> or FFM/L<sup>2</sup> at birth on growth and body composition over the subsequent two months further analyses were performed. When controlled for FM/L<sup>3</sup> Z-score at birth, higher IGF-I Z-scores at birth were associated with a reduction in FM/L<sup>3</sup> Z-score between birth and two months of age ( $R^2=0.01$ ,  $p=0.003$ ). This association was not seen with IGF-II Z-scores at birth ( $R^2=0.01$ ,  $p=0.07$ ). When controlling for FFM/L<sup>2</sup> Z-score at birth, there was no association between the change in FFM/L<sup>2</sup> Z-score over the first two months with IGF-I ( $R^2=0.01$ ,  $P=0.08$ ) or -II ( $R^2=0$ ,  $P=0.88$ ) Z-scores at birth.

## Discussion

This study demonstrated the relationships between IGF-I and -II concentrations at birth and neonatal body composition in a large well-defined cohort of healthy, term infants in Ireland. Lower IGF-I concentrations at birth were associated with reduced adiposity and lean mass, while lower IGF-II concentrations at birth were associated with reduced lean mass. Despite these significant associations, IGF concentrations accounted for a small amount of the variance in these body composition parameters. IGF-I and -II concentrations at birth were not associated with body composition parameters at two months. However lower IGF-I concentrations at birth were associated with a greater increase in both adiposity and lean body mass in both sexes between birth and two months.

The relationship between IGF-I concentration at birth and adiposity is not surprising. There are strong associations between IGF-I concentrations and birth weight(14, 28) and nutritional status(11). IGF-II is an important prenatal growth factor. Chorionic villi expression of *IGF2* mRNA is associated with birth weight(29) and paternally inherited *IGF2* mutations are associated with growth restriction(30). Animal models also support this important role of IGF-II in regulating prenatal growth(31). It is possible that IGF-II is more critical to growth prior to the third trimester(32), and this may explain the small association between body composition and IGF-II concentrations at birth.

Although IGF-I and -II concentrations at birth are not directly associated with body composition parameters at two months, there is a weak association with trajectory of accumulation of

adiposity. Significant changes occur in body composition during the first three months from birth, with a two-fold increase in FM corrected for length. Lean mass remains relatively stable during this time(16). The inverse relationship between IGF-I at birth and the change in adiposity may simply reflect the early “catch up” accumulation of body fat in infants with relative intrauterine growth restriction at birth. The same inverse association was seen between change in lean mass and IGF-I concentration at birth, indicating that this “catch-up” is not limited to adiposity but also occurs in lean mass. While significant, it should be noted that IGF-I concentrations accounted for only 5% of the variance in changes in FM/L<sup>3</sup> and over the first two months, and this reduced to 1% when corrected for FM/L<sup>3</sup> at birth. This suggests that IGF-I concentrations at birth only predict a small fraction of observed subsequent body composition changes.

The consequences of this relationship may extend to the interplay between intrauterine nutritional status, the GH/IGF axis and fetal programming of postnatal growth. Pregnancies complicated by limited access to nutrition are associated with increased IGF-I concentrations in adulthood, and it is hypothesized that chronic IGF suppression *in utero* results in an irreversible “overshoot” of this axis in later life(33). Children who have rapid early weight gain have increased BMI and insulin resistance as early as eight years of age(34), and SGA children with rapid early weight gain have persistent increases in adiposity when compared with similar weight, control children who were not born SGA(35). Although the mechanisms for these persistent metabolic complications are unknown, it is possible that low IGF-I concentration at birth is a biomarker for future risk.

Strengths of this study include the large, well-defined, relatively homogenous study population. All children were term, healthy and Caucasian, making confounding factors unlikely to contribute to these findings. However, this homogeneity may limit the generalizability of our findings to other populations. The large number of children with available body composition measurement is also unique for investigating these relationships. IGF-I and -II measurement by standard radioimmunoassays can be subject to assay interference by IGF binding proteins. However, LCMS was used in this study and this technique is not subject to interference from binding proteins(26, 27). A weakness of the study is the lack of available IGF-I and -II measurements at two months, to further investigate the hypothesis that IGF-I and -II may be useful biomarkers of metabolic risk during catch-up growth. Also, this study only included term infants and it is not known if the association is similar for preterm infants. Data were not available for the entire birth cohort, and it is not known if this introduced bias to our analysis.

We conclude that IGF-I concentrations at birth are associated with adiposity and lean mass at birth and the trajectory of FM accumulation over the first two months, although this only accounts for a small proportion of the variability seen in these parameters. Although IGF-I and -II play a significant role in prenatal growth, we have found that, in a cohort of healthy term infants, IGF-I and -II concentrations at birth are not major determinants of body composition changes over the first 2 months.

## References

1. Eriksson JG, Forsen T, Tuomilehto J, Osmond C, Barker DJ 2001 Early growth and coronary heart disease in later life: longitudinal study. *BMJ* 322:949-953.
2. Lopuhaa CE, Roseboom TJ, Osmond C, et al. 2000 Atopy, lung function, and obstructive airways disease after prenatal exposure to famine. *Thorax* 55:555-561.
3. Lussana F, Painter RC, Ocke MC, Buller HR, Bossuyt PM, Roseboom TJ 2008 Prenatal exposure to the Dutch famine is associated with a preference for fatty foods and a more atherogenic lipid profile. *Am J Clin Nutr* 88:1648-1652.
4. Neugebauer R, Hoek HW, Susser E 1999 Prenatal exposure to wartime famine and development of antisocial personality disorder in early adulthood. *JAMA* 282:455-462.
5. Hoek HW, Brown AS, Susser E 1998 The Dutch famine and schizophrenia spectrum disorders. *Soc Psychiatry Psychiatr Epidemiol* 33:373-379.
6. Gillman MW 2010 Early infancy as a critical period for development of obesity and related conditions. *Nestle Nutr Workshop Ser Pediatr Program* 65:13-20; discussion 20-14.
7. Ibanez L, Ong K, Dunger DB, de Zegher F 2006 Early development of adiposity and insulin resistance after catch-up weight gain in small-for-gestational-age children. *J Clin Endocrinol Metab* 91:2153-2158.
8. Ibanez L, Jimenez R, de Zegher F 2006 Early puberty-menarche after precocious pubarche: relation to prenatal growth. *Pediatrics* 117:117-121.
9. Stettler N, Zemel BS, Kumanyika S, Stallings VA 2002 Infant weight gain and childhood overweight status in a multicenter, cohort study. *Pediatrics* 109:194-199.



10. Abuzzahab MJ, Schneider A, Goddard A, et al. 2003 IGF-I receptor mutations resulting in intrauterine and postnatal growth retardation. *N Engl J Med* 349:2211-2222.
11. Hawkes CP, Grimberg A 2015 Insulin-like growth factor-I is a marker for the nutritional state. *Pediatr Endocrinol Rev* 13:465-477.
12. Wit JM, van Unen H 1992 Growth of infants with neonatal growth hormone deficiency. *Arch Dis Child* 67:920-924.
13. Giudice LC, de Zegher F, Gargosky SE, et al. 1995 Insulin-like growth factors and their binding proteins in the term and preterm human fetus and neonate with normal and extremes of intrauterine growth. *J Clin Endocrinol Metab* 80:1548-1555.
14. Hawkes CP, Murray DM, Kenny LC, et al. 2018 Correlation of Insulin-Like Growth Factor-I and -II Concentrations at Birth Measured by Mass Spectrometry and Growth from Birth to Two Months. *Horm Res Paediatr* 89:122-131.
15. Green LR, Kawagoe Y, Hill DJ, Richardson BS, Han VK 2000 The effect of intermittent umbilical cord occlusion on insulin-like growth factors and their binding proteins in preterm and near-term ovine fetuses. *J Endocrinol* 166:565-577.
16. Hawkes CP, Zemel BS, Kiely M, et al. 2016 Body Composition within the First 3 Months: Optimized Correction for Length and Correlation with BMI at 2 Years. *Horm Res Paediatr* 86:178-187.
17. Hawkes CP, Hourihane JO, Kenny LC, Irvine AD, Kiely M, Murray DM 2011 Gender- and Gestational Age-Specific Body Fat Percentage at Birth. *Pediatrics* 128:E645-E651.
18. Ibanez L, Sebastiani G, Lopez-Bermejo A, Diaz M, Gomez-Roig MD, de Zegher F 2008 Gender specificity of body adiposity and circulating adiponectin, visfatin, insulin, and

- insulin growth factor-I at term birth: relation to prenatal growth. *J Clin Endocrinol Metab* 93:2774-2778.
19. Chellakooty M, Juul A, Boisen KA, et al. 2006 A prospective study of serum insulin-like growth factor I (IGF-I) and IGF-binding protein-3 in 942 healthy infants: associations with birth weight, gender, growth velocity, and breastfeeding. *J Clin Endocrinol Metab* 91:820-826.
  20. O'Donovan SM, Murray DM, Hourihane JO, Kenny LC, Irvine AD, Kiely M 2014 Cohort profile: The Cork BASELINE Birth Cohort Study: Babies after SCOPE: Evaluating the Longitudinal Impact on Neurological and Nutritional Endpoints. *Int J Epidemiol* 44:764-775.
  21. Roggero P, Gianni ML, Amato O, Agosti M, Fumagalli M, Mosca F 2007 Measuring the body composition of preterm and term neonates: from research to clinical applications. *J Pediatr Gastroenterol Nutr* 45 Suppl 3:S159-162.
  22. Brozek J, Grande F, Anderson JT, Keys A 1963 Densitometric analysis of body composition: Revision of some quantitative assumptions. *Ann N Y Acad Sci* 110:113-140.
  23. ud Din Z, Emmett P, Steer C, Emond A 2013 Growth outcomes of weight faltering in infancy in ALSPAC. *Pediatrics* 131:e843-849.
  24. VanItallie TB, Yang MU, Heymsfield SB, Funk RC, Boileau RA 1990 Height-normalized indices of the body's fat-free mass and fat mass: potentially useful indicators of nutritional status. *Am J Clin Nutr* 52:953-959.
  25. Wells JC, Cole TJ, steam As 2002 Adjustment of fat-free mass and fat mass for height in children aged 8 y. *Int J Obes Relat Metab Disord* 26:947-952.

26. Bystrom C, Sheng S, Zhang K, Caulfield M, Clarke NJ, Reitz R 2012 Clinical utility of insulin-like growth factor 1 and 2; determination by high resolution mass spectrometry. *PLoS One* 7:e43457.
27. Bystrom CE, Sheng S, Clarke NJ 2011 Narrow mass extraction of time-of-flight data for quantitative analysis of proteins: determination of insulin-like growth factor-1. *Anal Chem* 83:9005-9010.
28. Christou H, Connors JM, Ziotopoulou M, et al. 2001 Cord blood leptin and insulin-like growth factor levels are independent predictors of fetal growth. *J Clin Endocrinol Metab* 86:935-938.
29. Demetriou C, Abu-Amero S, Thomas AC, et al. 2014 Paternally expressed, imprinted insulin-like growth factor-2 in chorionic villi correlates significantly with birth weight. *PLoS One* 9:e85454.
30. Begemann M, Zirn B, Santen G, et al. 2015 Paternally Inherited IGF2 Mutation and Growth Restriction. *N Engl J Med* 373:349-356.
31. DeChiara TM, Efstratiadis A, Robertson EJ 1990 A growth-deficiency phenotype in heterozygous mice carrying an insulin-like growth factor II gene disrupted by targeting. *Nature* 345:78-80.
32. Langford K, Nicolaides K, Miell JP 1998 Maternal and fetal insulin-like growth factors and their binding proteins in the second and third trimesters of human pregnancy. *Hum Reprod* 13:1389-1393.
33. Elias SG, Keinan-Boker L, Peeters PH, et al. 2004 Long term consequences of the 1944-1945 Dutch famine on the insulin-like growth factor axis. *Int J Cancer* 108:628-630.

34. Ong KK, Petry CJ, Emmett PM, et al. 2004 Insulin sensitivity and secretion in normal children related to size at birth, postnatal growth, and plasma insulin-like growth factor-I levels. *Diabetologia* 47:1064-1070.
35. Ibanez L, Lopez-Bermejo A, Diaz M, de Zegher F 2011 Catch-up growth in girls born small for gestational age precedes childhood progression to high adiposity. *Fertil Steril* 96:220-223.

Author accepted manuscript

## Figure Legends

**Figure 1:** The number of subjects recruited who 1) were born at term, 2) had umbilical cord blood samples available and 3) had body composition measurement performed at birth and two months.

**Figure 2:** Scatter-plot and linear regression comparing IGF-I (A and B) and -II (C and D) concentrations with age- and sex-corrected FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at birth. The associations between IGF-I concentration and FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores and the association between IGF-II concentration and FFM/L<sup>2</sup> Z-score at birth were significant.

**Figure 3:** Scatter-plot and linear regression comparing IGF-I and -II concentrations at birth with age- and sex-corrected FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at 2 months.

**Table 1:** Characteristics of population. Unless otherwise stated, mean (SD) are presented.

	<b>Males</b>	<b>Females</b>	<b>All</b>
Number, n	317	284	601
Gestational age, weeks	40.2 (1.1)	40.2 (1)	40.2 (1.1)
Birth weight, kg	3.57 (0.45)	3.49 (0.45)	3.53 (0.45)
Weight at 2 months, kg	5.66 (0.68)	5.3 (0.57)	5.49 (0.66)
Length at birth, m	0.51 (0.02)	0.50 (0.02)	0.50 (0.02)
Length at 2 months, m	0.59 (0.02)	0.58 (0.02)	0.59 (0.02)
IGF-I, ng/ml	49.6 (23.9)	56.6 (24.3)	52.9 (24.3)
IGF-I, Z-score	0.07 (1)	-0.004 (1)	0.03 (1)
IGF-II, ng/ml	421.4 (98.1)	428.5 (103.7)	424.8 (100.8)
IGF-II, Z-score	0 (1)	-0.01 (1)	- 0.004 (1)
FM/L <sup>3</sup> at birth, kg/m <sup>3</sup>	2.7 (1.2)	3.2 (1.2)	2.9 (1.2)
FM/L <sup>3</sup> at birth, Z-score	0 (1)	-0.02 (1)	-0.01 (1)
FM/L <sup>3</sup> at 2m, kg/m <sup>3</sup>	5.9 (1.5)	6.2 (1.4)	6 (1.5)
FM/L <sup>3</sup> at 2m, Z-score	0 (1)	0.09 (1)	0.04 (1)
FFM/L <sup>2</sup> at birth, kg/m <sup>2</sup>	11.8 (0.9)	11.5 (0.9)	11.7 (0.9)
FFM/L <sup>2</sup> at birth, Z-score	0 (1)	0.06 (1)	0.02 (1)
FFM/L <sup>2</sup> at 2m, kg/m <sup>2</sup>	12.8 (0.8)	12.1 (0.8)	12.5 (0.9)
FFM/L <sup>2</sup> at 2m, Z-score	0.04 (1)	0.04 (1)	0.04 (1)
Change in FM/L <sup>3</sup> Z-score from birth to 2-months	0 (1.2)	0.1 (1.2)	0.05 (1.2)
Change in FFM/L <sup>2</sup> Z-score from birth to 2-months	0.04 (1)	-0.02 (1.1)	0.01 (1)

**Table 2:** The relationship between sex- and gestational age-corrected FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at birth and two months, with sex- and gestational age-corrected IGF-I and IGF-II Z-scores at birth. For this analysis, IGF-I and -II Z-scores are the dependent variables and FM/L<sup>3</sup> or FFM/L<sup>2</sup> measures are independent variables.

\*Partial correlation controlling for FM/L<sup>3</sup> Z-Score at birth, \*\* Partial correlation controlling for FFM/L<sup>2</sup> Z-Score at birth

	IGF-I Z-score			IGF-II Z-score		
	R <sup>2</sup>	Regression coefficient (SEM)	P	R <sup>2</sup>	Regression coefficient (SEM)	P
FM/L <sup>3</sup> Z-score at birth	0.05	0.23 (0.04)	<0.001	0.001	0.04 (0.04)	0.4
FM/L <sup>3</sup> Z-score at 2 months	0.002	-0.05 (0.04)	0.3	0.004	-0.06 (0.04)	0.1
FFM/L <sup>2</sup> Z-score at birth	0.04	0.2 (0.04)	<0.001	0.01	0.1 (0.04)	0.02
FFM/L <sup>2</sup> Z-score at 2 months	0.001	0.03 (0.04)	0.5	0.002	0.04 (0.04)	0.3
Change in FM/L <sup>3</sup> Z-score from birth to 2 months*	0.01	-0.12 (0.04)	0.003	0.01	-0.07 (0.04)	0.07
Change in FFM/L <sup>2</sup> Z-score from birth to 2 months**	0.01	-0.07 (0.04)	0.08	0	-0.01 (0.04)	0.88




Figure 1: The number of subjects recruited who 1) were born at term, 2) had umbilical cord blood samples available and 3) had body composition measurement performed at birth and two months.

66x37mm (300 x 300 DPI)






Figure 2: Scatter-plot and linear regression comparing IGF-I (A and B) and -II (C and D) concentrations with age- and sex-corrected FM/L3 and FFM/L2 Z-scores at birth. The associations between IGF-I concentration and FM/L3 and FFM/L2 Z-scores and the association between IGF-II concentration and FFM/L2 Z-score at birth were significant.

110x114mm (300 x 300 DPI)




Figure 3: Scatter-plot and linear regression comparing IGF-I and -II concentrations at birth with age- and sex-corrected FM/L<sup>3</sup> and FFM/L<sup>2</sup> Z-scores at 2 months.

110x118mm (300 x 300 DPI)